

ADOPTED: dd mmmm 2017

doi:10.2903/j.efsa.2017.NNNN

Draft Scientific Opinion on the safety and suitability for use by infants of follow-on formulae with a protein content of at least 1.6 g/100 kcal

EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA),
Dominique Turck, Jean-Louis Bresson, Barbara Burlingame, Tara Dean, Susan Fairweather-Tait, Marina Heinonen, Karen Ildico Hirsch-Ernst, Inge Mangelsdorf, Harry J McArdle, Androniki Naska, Monika Neuhäuser-Berthold, Grażyna Nowicka, Kristina Pentieva, Yolanda Sanz, Anders Sjödin, Martin Stern, Daniel Tomé, Henk Van Loveren, Marco Vinceti, Peter Willatts, Mary Fewtrell, Hildegard Przyrembel

Abstract

Following a request from the European Commission, the EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA) was asked to deliver a scientific opinion on the safety and suitability for use by infants of follow-on formulae (FOF) based on cow's milk intact protein with a protein content of at least 1.6 g/100 kcal (rounded value) that meet otherwise the requirements of relevant EU legislation. If the formula under evaluation is considered to be safe and suitable for use by infants, the NDA Panel is also asked to advise on whether FOF based on goat's milk intact protein, soy protein isolates or protein hydrolysates are also safe and suitable for infants under the same conditions. The Panel concludes that the use of FOF with a protein content of at least 1.6 g/100 kcal from either intact cow's milk protein or intact goat's milk protein otherwise complying with the requirements of relevant EU legislation is safe and suitable for infants living in Europe with access to complementary foods of a sufficient quality. This conclusion does not apply to IF. The Panel also concludes that the safety and suitability of FOF with a protein content of at least 1.6 g/100 kcal manufactured from either protein hydrolysates or soy protein isolates cannot be established with the available data. The same conclusion applies to IF. The NDA Panel endorsed this draft scientific opinion on 14 December 2016 for public consultation, to which stakeholders are encouraged to contribute. The document will be revised and updated according to the comments received, where appropriate.

© 2017 European Food Safety Authority. *EFSA Journal* published by John Wiley and Sons Ltd on behalf of European Food Safety Authority.

Keywords: protein, infants, follow-on formula, safety, suitability, growth

Requestor: European Commission

Question number: EFSA-Q-2016-00275

Correspondence: nda@efsa.europa.eu

37 **Panel members:** Jean-Louis Bresson, Barbara Burlingame, Tara Dean, Susan Fairweather-Tait,
38 Marina Heinonen, Karen Ildico Hirsch-Ernst, Inge Mangelsdorf, Harry J McArdle, Androniki Naska,
39 Monika Neuhäuser-Berthold, Grażyna Nowicka, Kristina Pentieva, Yolanda Sanz, Alfonso Siani, Anders
40 Sjödin, Martin Stern, Daniel Tomé, Dominique Turck, Henk Van Loveren, Marco Vinceti and Peter
41 Willatts.

42 **Acknowledgements:** The Panel wishes to thank the members of the Working Group on Infant
43 Nutrition: Jean-Louis Bresson, Mary Fewtrell, Hildegard Przyrembel, Dominique Turck, and the
44 external experts: Rosangela Marchelli and Daniel Tomé for the preparatory work on this scientific
45 output, and EFSA staff members: Davide Arcella, Krizia Ferrini, Ariane Titz and Silvia Valtueña
46 Martínez for the support provided to this scientific output.

47 **Suggested citation:** EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies),
48 2017. Scientific Opinion on the safety and suitability for use by infants of follow-on formulae with a
49 protein content of at least 1.6 g/100 kcal. EFSA Journal 2017;volume(issue):NNNN, 31 pp.
50 doi:10.2903/j.efsa.2017.NNNN

51 **ISSN:** 1831-4732

52 © 2017 European Food Safety Authority. *EFSA Journal* published by John Wiley and Sons Ltd on
53 behalf of European Food Safety Authority.

54 This is an open access article under the terms of the [Creative Commons Attribution-NoDerivs](https://creativecommons.org/licenses/by/4.0/) License,
55 which permits use and distribution in any medium, provided the original work is properly cited and no
56 modifications or adaptations are made.

57



The EFSA Journal is a publication of the European Food Safety Authority, an agency of the European Union.



58

59 Summary

60 Following a request from the European Commission, the EFSA Panel on Dietetic Products, Nutrition
61 and Allergies (NDA) was asked to deliver a scientific opinion on the safety and suitability for use by
62 infants of follow-on formulae (FOF) based on cow's milk intact protein with a protein content of at
63 least 1.6 g/100 kcal (rounded value) that meet otherwise the requirements of relevant EU legislation.
64 If the formula under evaluation is considered to be safe and suitable for use by infants, the NDA Panel
65 is also asked to advise on whether FOF based on goat's milk intact protein, soy protein isolates or
66 protein hydrolysates are also safe and suitable for infants under the same conditions.

67 For the scientific assessment, the NDA Panel considered: a) the dietary protein requirements of
68 infants in the second half of the first year of life, b) the protein content of breast milk during the first
69 year of lactation, c) dietary protein intake of infants in Europe from breast milk, formula and
70 complementary food (CF), d) the overall contribution that a FOF with a protein content of 1.6 g/100
71 kcal could make towards protein requirements in the target population, assuming access to CF of a
72 sufficient quality, following established feeding guidelines in Europe (e.g. from Member States), and e)
73 the application submitted by the food business operator, including two intervention studies in healthy
74 term infants.

75 The Panel notes that:

- 76 a) Population Reference Intakes (PRIs) of 10 g protein per day for girls and 11 g protein per day
77 for boys aged 6 months and a PRI of 11 g protein per day for girls and 12 g protein per day
78 for boys aged 12 months have been established,
79
- 80 b) The mean content of true protein in breast milk by the end of the third month of lactation
81 ranges between 1.3 and 1.6 g/100kcal, tends to decrease thereafter to about 1.1-1.4
82 g/100kcal by the end of the fourth month, and tends to remain fairly stable thereafter,
83
- 84 c) The P5th and P2.5th of total protein intake in non-breastfed infants aged 6 to 12 months living
85 in Europe was around or above the PRI for protein for that age group in all the studies and
86 surveys available,
87
- 88 d) The P5th and P2.5th of total protein intake resulting from the consumption of FOF with a
89 protein content of 1.6 g/100kcal would remain at about or above the PRI for protein for
90 infants aged 6 to 12 months who are not breastfed, and
91
- 92 e) The two randomised, double-blind, controlled intervention studies provided by the applicant
93 showed no differences in growth patterns between healthy term infants who consumed
94 formulae with protein contents of 1.61 g/100 kcal and 1.65 g/100 kcal from three months of
95 age onwards and infants who consumed formulae with protein contents of 2.15 g/100 kcal
96 and 2.70 g/100 kcal, respectively. The control formula used in these studies contained 0.35
97 g/100 kcal (US study) and 0.90 g/100 kcal (Chile study) more protein than the current
98 minimum requirement for protein content of a FOF (1.8 g/100 kcal).

99 The Panel also notes that the studies submitted were not specifically designed to meet the regulatory
100 definitions for either IF or FOF laid down in Regulation (EU) No 609/2013¹, and that the information
101 provided in relation to the type and amount of CF was not sufficient to calculate total energy and
102 protein intake, nor the relative contribution of formulae and CF to total energy and protein intake.
103 Therefore, the Panel considers that these studies do not provide, on their own, sufficient information
104 to conclude on the safety and suitability of a FOF with a protein content of 1.6 g protein/100kcal.

105 The Panel notes, however, that:

¹ Regulation (EU) No 609/2013 of the European Parliament and of the Council of 12 June 2013 on food intended for infants and young children, food for special medical purposes, and total diet replacement for weight control and repealing Council Directive 92/52/EEC, Commission Directives 96/8/EC, 1999/21/EC, 2006/125/EC and 2006/141/EC, Directive 2009/39/EC of the European Parliament and of the Council and Commission Regulations (EC) No 41/2009 and (EC) No 953/2009, OJ L 181, 29.6.2013, p. 35–56

- 106 a) the protein content of human milk tends to decrease with feeding time to about 1.1-1.4
107 g/100kcal by the end of the fourth month of lactation, remaining fairly stable thereafter,
- 108 b) that P5th and P2.5th of protein intake from all sources (breast milk, formula and CF) in
109 European infants between 6 and 12 months of age are at or above the PRI for protein for
110 that age group.
- 111 c) P5th and P2.5th of protein intake from all sources (formula and CF) in European infants
112 between 6 and 12 months of age who are not breastfed would remain at or above the PRI
113 for protein for that age group by assuming a protein content of 1.6 g/100kcal in all FOF.
- 114 d) the two human intervention studies provided by the applicant did not show an adverse
115 impact on growth resulting from the use of a formula containing about 1.6 g of
116 protein/100kcal as compared to control formulae containing 2.15 or 2.70 g of protein/100
117 kcal or the breastfed reference group.

118 Therefore, the Panel concludes that the use of FOF with a protein content of at least 1.6 g/100
119 kcal from intact cow's milk protein otherwise complying with the requirements of relevant EU
120 legislation is safe and suitable for infants living in Europe with access to complementary foods of
121 a sufficient quality. This conclusion does not apply to IF.

122 On the basis of:

- 123 a) a previous evaluation by the Panel on the safety and suitability of goat's milk protein as a
124 source of protein in IF and FOF (EFSA NDA Panel, 2012b), and
- 125 b) the Panel's conclusions regarding the safety and suitability of FOF with a protein content of at
126 least 1.6 g/100 kcal from intact cow's milk protein otherwise complying with the
127 requirements of relevant EU legislation,

128 the Panel concludes that the use of FOF with a protein content of at least 1.6 g/100 kcal from intact
129 goat's milk protein otherwise complying with the requirements of relevant EU legislation is safe and
130 suitable for infants living in Europe with access to complementary foods of a sufficient quality. This
131 conclusion does not apply to IF.

132 The Panel considers, however, that the safety and suitability of each FOF (and IF) manufactured from
133 protein hydrolysates have to be established by clinical evaluation in the target population (EFSA NDA
134 Panel, 2014). The Panel also considers that, given the higher minimum protein requirements
135 established for FOF (and IF) manufactured from soy protein isolates (i.e. 2.25 g/100 kcal) and the
136 lack of data available on the use of FOF from soy protein isolates in the target population, additional
137 studies are required to establish the safety and suitability of FOF manufactured from soy protein
138 isolates with a protein content of at least 1.6 g/100 kcal. Therefore, the Panel concludes that the
139 safety and suitability of FOF with a protein content of at least 1.6 g/100 kcal manufactured from either
140 protein hydrolysates or soy protein isolates cannot be established with the available data. The same
141 conclusion applies to IF.

142

143 **Table of contents**

144		
145	Abstract.....	1
146	Summary.....	3
147	1. Introduction.....	6
148	1.1. Background and Terms of Reference as provided by the European Commission.....	6
149	1.1.1. Background.....	6
150	1.1.2. Terms of Reference.....	6
151	1.2. Interpretation of the Terms of Reference.....	6
152	2. Data and Methodologies.....	7
153	2.1. Data.....	7
154	2.2. Methodologies.....	7
155	3. Assessment.....	8
156	3.1. Dietary protein requirements of infants in the second half of the first year of life.....	8
157	3.2. Protein content of breast milk during the first year of lactation.....	8
158	3.3. Dietary protein intake of infants in Europe.....	10
159	3.4. Contribution that a FOF with a protein content of at least 1.6 g/100 kcal could make towards protein requirements in the target population.....	18
161	3.5. Application submitted by the food business operator.....	19
162	3.5.1. Composition of the formulae used in the two human intervention studies.....	19
163	3.5.2. Human intervention studies.....	20
164	3.6. Comparison between the human intervention studies provided and European dietary surveys with respect to mean energy and protein intake from formula and complementary food in the target population.....	22
167	4. Conclusions.....	23
168	4.1. On the safety and suitability for use by infants of FOF with a protein content of at least 1.6 g/100 kcal from intact cow's milk protein otherwise complying with the requirements of relevant EU legislation.....	23
171	4.2. On the safety and suitability for use by infants of FOF with a protein content of at least 1.6 g/100 kcal from goat's milk intact protein, soy protein isolates or protein hydrolysates otherwise complying with the requirements of relevant EU legislation.....	23
174	Documentation provided to EFSA.....	25
175	References.....	25
176	Abbreviations.....	27
177	Appendix A – Absolute weight, weight gain and weight change in the "US study".....	28
178	Appendix B – Length and head circumference in the "US study".....	29
179	Appendix C – Absolute weight, weight gain and weight change in the "Chile study".....	30
180	Appendix D – Length and head circumference in the "Chile study".....	31
181		

182

183 **1. Introduction**

184 **1.1. Background and Terms of Reference as provided by the European** 185 **Commission**

186 **1.1.1. Background**

187 Commission Directive 2006/141/EC² lays down requirements for infant formulae and follow-on
188 formulae placed on the market in the EU. Among others, it establishes that follow-on formula
189 manufactured from cows' milk intact protein shall contain at least 1.8 g protein/100 kcal (Annex II,
190 point 2.1).

191 Commission delegated Regulation (EU) 2016/127³ revises the rules of Commission Directive
192 2006/141/EC and shall repeal and replace the Directive from 22 February 2020. Annex II, point 2.1 of
193 the delegated Regulation maintains the minimum protein content of follow-on formula manufactured
194 from cows' milk intact protein at 1.8 g/100 kcal.

195 The Commission has received a request from a food business operator for placing on the market a
196 follow-on formula based on cow's milk intact protein with a protein content of at least 1.61 g/100 kcal,
197 which is below the permitted levels of Directive 2006/141/EC and delegated Regulation (EU)
198 2016/127. In order to consider such request, the Commission needs to obtain the advice of the
199 European Food Safety Authority and has asked the food business operator to send the scientific
200 dossier to the Authority for assessment.

201 **1.1.2. Terms of Reference**

202 In accordance with Article 29 of Regulation (EC) No 178/2002⁴, the European Commission requests
203 the European Food Safety Authority to issue an opinion on the safety and suitability for use by infants
204 of a follow-on formula based on cow's milk intact protein with a protein content of at least 1.61 g/100
205 kcal.

206 If the formula under evaluation is considered to be safe and suitable for use by infants, the European
207 Food Safety Authority is asked to advise whether a level of at least 1.61 g protein/100 kcal would be
208 applicable to all follow-on formulae. If this is not the case, the Authority is asked to advise on the
209 specific criteria that need to be satisfied for the safety and suitability of such formulae to be
210 demonstrated.

211 **1.2. Interpretation of the Terms of Reference**

212 The Panel interprets the terms of reference provided by the European Commission in the context of
213 the background information given and the application submitted. The Panel understands that the
214 European Commission seeks advice on:

215 (a) whether a follow-on formula (FOF) based on cow's milk intact protein with a minimum protein
216 content of 1.6 g/100 kcal (rounded value) is safe and suitable for infants provided that it meets
217 otherwise the requirements⁵ of relevant EU legislation⁶, and if so

218 (b) whether FOF based on goat's milk intact protein, soy protein isolates or protein hydrolysates with
219 a minimum protein content of 1.6 g/100 kcal (rounded value) are also safe and suitable for infants
220 provided that they meet otherwise the requirements of relevant EU legislation.

² Commission Directive 2006/141/EC of 22 December 2006 on infant formulae and follow-on formulae and amending Directive 1999/21/EC, OJ L 401, 30.12.2006, p. 1

³ Commission Delegated Regulation (EU) 2016/127 of 25 September 2015 supplementing Regulation (EU) No 609/2013 of the European Parliament and of the Council as regards the specific compositional and information requirements for infant formula and follow-on formula and as regards requirements on information relating to infant and young child feeding, OJ L 25, 2.2.2016, p. 1

⁴ Regulation (EC) No 178/2002 of the European Parliament and of the Council of 28 January 2002 laying down the general principles and requirements of food law, establishing the European Food Safety Authority and laying down procedures in matters of food safety, OJ L 31, 1.2.2002, p. 1

⁵ Including the requirements with respect to the amino acid profile

⁶ Directive 2006/141/EC to be replaced by delegated Regulation (EU) 2016/127

221 **2. Data and Methodologies**

222 **2.1. Data**

223 EFSA was provided with a dossier related to a FOF based on cow's milk intact protein containing a
224 minimum of 1.61 g protein per 100 kcal but otherwise complying with the compositional criteria laid
225 down in Directive 2006/141/EC. The dossier includes two intervention studies in infants, named "US
226 study" (Hayes and Northington, 2014, unpublished study report #1; published as Ziegler et al., 2015),
227 and "Chile study" (Yao, 2014, unpublished study report #2; published as Inostroza et al., 2014). The
228 dossier was supplemented, upon request of EFSA, with additional information provided by the
229 applicant on 28 June 2016 and on 29 September 2016. The intervention studies provided in the
230 dossier were designed to assess the growth pattern of infants receiving a formula with standard
231 protein content for the first three months of life and thereafter a formula with a protein content which
232 is lower than currently authorised. These studies aimed to investigate whether lower protein content
233 in formula to be fed from three to 12 months of age, in line with the decrease in the protein content
234 of breast milk during that feeding period, would lead to growth rates closer to those of breast-fed
235 infants, as compared to infants fed a "standard" formula.

236 The Panel will also take into account in the current assessment its previous opinions on Dietary
237 Reference Values for protein (EFSA NDA Panel, 2012a), on nutrient requirements and dietary intake of
238 infants and young children in Europe (EFSA NDA Panel, 2013), and on the essential composition of IF
239 and FOF (EFSA NDA Panel, 2014), as well as data on the protein content of breast milk.

240 **2.2. Methodologies**

241 As outlined in the Panel's previous opinion on the essential composition of IF and FOF (EFSA NDA
242 Panel, 2014), the minimum amounts of nutrients in formulae, including protein, should be based on
243 generally accepted scientific evidence. While for IF compositional requirements may be based on the
244 energy and nutrient requirements of infants and on the results of intervention studies in the target
245 population in which the formula is the only source of energy and nutrients, evidence for proposing
246 compositional requirements for foods which are not the sole source of energy and nutrients, such as
247 FOF, is less strong, as other foods contribute to nutrient and energy intake in variable amounts. In its
248 previous opinion, when proposing compositional requirements for FOF, the Panel assumed that
249 complementary food (CF) would compensate for the higher energy and nutrient requirements of older
250 infants and for the lower formula intake during that period. This is based on the assumption that
251 infants in the target population have access to CF of a sufficient quality, following established feeding
252 guidelines in Europe (e.g. from Member States).

253 For the present assessment of whether a FOF based on cow's milk intact protein with a protein
254 content of at least 1.6 g/100 kcal (rounded value) is safe and suitable for infants provided that it
255 meets otherwise the requirements⁷ of relevant EU legislation⁸, the Panel will consider:

- 256 a) Dietary protein requirements of infants in the second half of the first year of life;
- 257 b) Protein content in breast milk during the first year of lactation;
- 258 c) Dietary protein intake of infants in Europe from breast milk, formula and CF;
- 259 d) The overall contribution that a FOF with a protein content of 1.6 g/100 kcal could make
260 towards protein requirements in the target population, assuming access to CF of a sufficient
261 quality, following established feeding guidelines in Europe (e.g. from Member States);
- 262 e) The application submitted by the food business operator, including two intervention studies in
263 healthy term infants.

264 The evaluation of the intervention studies provided by the food business operator will follow the
265 general principles for the assessment of a modification of the composition of IFs or FOFs outside the
266 established standards as laid down by the Scientific Committee on Food (SCF) (SCF, 2003). In
267 addition, the recommendations for the assessment of the safety and suitability of formulae for term

⁷ Including the requirements with respect to the amino acid profile

⁸ Directive 2006/141/EC to be replaced by delegated Regulation (EU) 2016/127

268 infants of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition (ESPGHAN)
269 (Aggett et al., 2001), of the Committee on the Evaluation of the Addition of Ingredients New to Infant
270 Formula of the Food and Nutrition Board of the United States (US) Institute of Medicine (IoM, 2004)
271 and of the American Academy of Pediatrics (AAP, 1988), will be taken into account.

272 **3. Assessment**

273 **3.1. Dietary protein requirements of infants in the second half of the** 274 **first year of life**

275 Dietary protein is an essential component of the diet, supplying the body with nitrogen (N) and amino
276 acids as well as other non-protein metabolically active nitrogenous substances. The protein
277 requirement of infants and young children comprises two components, the maintenance requirement
278 and the growth requirement (EFSA NDA Panel, 2012a). In its previous opinion, the Panel established
279 an average maintenance requirement of 0.66 g protein/kg body weight per day (105 mg N/kg body
280 weight per day) for infants and young children aged from 6 to < 36 months, which was derived from
281 nitrogen balance studies in adults. The average protein requirement for growth was estimated from
282 average daily rates of protein deposition calculated from studies on whole-body potassium deposition,
283 and adjusted by an efficiency of utilisation of dietary protein for growth of 58 %. Together, these
284 amounts constitute an AR, to which 1.96 standard deviations were added to derive a PRI. Thus, a PRI
285 of 10 g protein per day for girls and 11 g protein per day for boys aged 6 months and a PRI of 11 g
286 protein per day for girls and 12 g protein per day for boys aged 12 months were established.

287 **3.2. Protein content of breast milk during the first year of lactation**

288 Estimating the true protein content of breast milk is challenging because of the non-protein nitrogen
289 fraction contained in it. Total nitrogen in human milk represents both protein, about 75%, and non-
290 protein nitrogen, which is made up of urea (up to 50% of the non-protein nitrogen), amino acids and
291 other nitrogenous compounds (SCF, 2003; WHO/FAO/UNU, 2007). The amount of nitrogen used by
292 infants for protein synthesis is likely to include that from true protein, free amino acids and small
293 peptides, and a proportion of urea nitrogen. Therefore, the amount of nitrogen in breast milk used for
294 protein synthesis by infants is between the true protein content and the crude protein calculated from
295 total nitrogen.

296 A meta-analysis of 41 published studies reporting on pre-term (26 studies, 843 mothers) and term (30
297 studies, 2299 mothers) breast milk composition is available (Gidrewicz and Fenton, 2014). Energy was
298 estimated in 11 studies using bomb calorimetry, and in five studies by calculation using values for the
299 energy contributions from fat, protein, and carbohydrate. Protein was estimated based on total
300 nitrogen in 23 studies and as a true protein estimate in 15 studies. Data on mean energy and protein
301 content of breast milk from mothers of term infants by week of lactation is shown in **Table 1**.

302

303 **Table 1:** Mean energy and protein content of breast milk from mothers of term infants by week
 304 of lactation¹

Time	Mean energy (SD) (kcal/100mL)		Mean protein (SD) (g/100mL)		Mean protein (SD) (g/100kcal)	
	Bomb calorimetry	Calculated	True protein	Protein calculated from total nitrogen	True protein	Protein calculated from total nitrogen
4-7 d	66 (9)	68 (10)	1.6 (0.3)	2.0 (0.5)	2.4 (0.5)	2.9 (0.7) -3.0 (0.8)
2 wk	66 (9)	-	1.3 (0.2)	1.8 (0.4)	2.0 (0.3)	2.7 (0.63)
3-4 wk	66 (8)	70 (9)	1.1 (0.2)	1.5 (0.3)	1.6 (0.3) -1.7 (0.3)	2.1 (0.4) -2.3 (0.5)
5-6 wk	63 (7)	-	1.0 (0.1)	1.1 (0.2)	1.6 (0.2)	1.7 (0.3)
7-9 wk	63 (7)	69 (10)	0.9 (0.1)	1.3 (0.2)	1.3-1.4 (0.2)	2.1 (0.3)
10-12 wk	63 (8)	68 (9)	1.0 (0.1)	1.2 (0.2)	1.5-1.6 (0.2)	1.8 (0.3) -1.9 (0.3)

305 ¹Adapted from Gidrewicz and Fenton (2014)

306 The true protein content of breast milk gradually decreased from the first week of lactation, being
 307 about 1.0 g/100mL (corresponding to about 1.8-1.9 g/100kcal) by the third month.

308 **Table 2** shows the energy (calculated) and the interval from birth according to the true protein
 309 content of breast milk samples (n=2554) donated by 224 mothers of mostly term infants to a milk
 310 bank in Denmark (Michaelsen et al., 1990). The mean true protein content of all samples combined
 311 was 0.9 g/100mL.

312 **Table 2:** Time interval from birth according to the protein content of breast milk¹

True protein (g/100mL)	Mean interval from birth (weeks)	Mean (SD) energy (kcal/100mL)	True protein (g/100kcal)	No. of samples
≥ 1.3	3-4	74.7 (9.8)	≥ 1.7	70
1.1 – 1.29	6-7	71.4 (9.8)	1.5-1.8	193
0.9 – 1.09	11-12	67.4 (9.8)	1.3-1.6	572
0.7 – 0.89	15-16	64.6 (9.8)	1.1-1.4	800
< 0.7	19-20	64.2 (9.3)	< 1.1	108

313 ¹Adapted from Michaelsen et al. (1990)

314 The mean interval from birth gradually decreased with the increasing content of protein in breast milk.
 315 It was 3-4 weeks for samples containing ≥1.3 g/100mL and 19-20 weeks for samples containing <0.7
 316 g/100mL. The mean interval from birth was 11-12 weeks for samples containing 0.9-1.09 g/100mL,
 317 which is consistent with the protein content of breast milk by the third month of lactation (1.0
 318 g/100mL) reported by Gidrewicz and Fenton (2014).

319 Other studies have reported on the protein content of breast milk during the first year of lactation,
 320 although the number of breast milk samples analyses in those studies is low (Allen et al., 1991;
 321 Nommsen et al., 1991; Mitoulas et al., 2002).

322 In the context of the DARLING study, Nommsen et al. (1991) assessed the composition of breast milk
 323 in samples taken at 3, 6, 9 and 12 months of lactation in healthy mothers of term infants. The gross
 324 energy content and the total protein content of the breast milk samples are given in **Table 3**. Protein
 325 was analysed using a modified Lowry assay with bovine serum albumin as the standard, a method
 326 which tends to result in slightly elevated values for total protein (Nommsen et al., 1991).

327

328 **Table 3:** Gross energy and total protein content of breast milk during the first year of lactation¹

Month of lactation	n	Mean (SD) gross energy (kcal/100mL)	Mean (SD) total protein (g/100mL)	Mean (SD) total protein (g/100kcal)
3	58	697 (97)	1.2 (0.2)	1.7 (1.5)
6	45	707 (92)	1.1 (0.2)	1.6 (1.6)
9	28	709 (74)	1.2 (0.8)	1.6 (10.8)
12	21	706 (110)	1.2 (0.2)	1.7 (1.3)

329 ¹Adapted from Nommsen et al. (1991)

330 The mean total protein at three months of lactation is comparable to the protein content calculated
 331 from total nitrogen (1.2 g/100mL, 1.8 g/100kcal) for that time period in the meta-analysis by
 332 Gidrewicz and Fenton (2014), and remained fairly stable until the twelfth month of lactation, which is
 333 in agreement with what has been reported in other studies (Allen et al., 1991; Mitoulas et al., 2002).

334 The Panel notes that the mean content of true protein in breast milk by the end of the third month of
 335 lactation ranges between 1.3 and 1.6 g/100kcal, tends to decrease thereafter to about 1.1-1.4
 336 g/100kcal by the end of the fourth month, and tends to remain fairly stable thereafter.

337 3.3. Dietary protein intake of infants in Europe

338 Data on mean energy and protein intake in infants living in Europe were gathered from published
 339 studies (**Table 4**). Details about the dietary data collection and on the assessment of breast milk
 340 intake are given in **Table 5**.

341 Data on mean energy and protein intake in infants living in Europe were also gathered from dietary
 342 surveys for which sufficient data were available in the EFSA Comprehensive European Food
 343 Consumption Database (**Table 6**)⁹.

344 From the dietary surveys available in the EFSA Comprehensive European Food Consumption
 345 Database, mean energy and protein intake from formula and from CF in non-breastfed infants, and
 346 from CF only in (exclusively or partially) breastfed infants were also calculated (**Table 7**). Mean
 347 energy and protein intake by food group in non-breastfed infants are given in **Tables 8** (for infants 4
 348 to <6 months of age) and **9** (for infants aged ≥6 to 12 months).

⁹ Details about the dietary surveys included in the EFSA Comprehensive Database are available at: <https://dwh.efsa.europa.eu/bi/asp/Main.aspx?rwtrep=001>

Table 4: Mean energy and protein intake in infants living in Europe from published studies

Age	Country	Study	Breastfeeding ¹ (%)	N	Mean energy (SD) ² (kcal/d) ³	Mean protein (SD) ² (g/d)	P5 th of protein intake ³	P2.5 th of protein intake ⁴	Mean E% as protein ²
4 mo	UK	ALSPAC	All males	262	658 (123)	15.7 (3.4)	10.1	9.0	9.5
			All females	214	604 (118)	14.5 (3.3)	9.04	8.0	9.6
			100, no solids	53	626 (-)	12 (-)	-	-	7.6
			0, no solids	42	583 (-)	13 (-)	-	-	8.9
			100, solids	209	646 (-)	13 (-)	-	-	8.1
			0, solids	441	640 (-)	15 (-)	-	-	9.4
			Mixed, solids	107	667 (-)	14 (-)	-	-	8.4
6 mo	France	National	0	58	668	20.0 (5.60)	10.7	-	11.9
	UK	Southampton	28	50	709 (652-818) ⁵	19.7 (17.4-23.6) ⁵	-	-	11.1 ⁷
	Germany	DONALD	48	302	645 (119)	16.3 (5.4)	7.4	5.7	10.1
	Belgium	CHOP	0	746	679 (659-702) ⁶	19 (17-20) ⁶	-	-	11.2
	Germany				634 (615-655) ⁶	19 (18-20) ⁶	-	-	12.0
	Italy				731 (710-753) ⁶	23 (22-24) ⁶	-	-	12.6
	Poland				725 (702-743) ⁶	21 (20-22) ⁶	-	-	11.6
	Spain				736 (719-754) ⁶	18 (18-19) ⁶	-	-	10.0
8 mo	Belgium	CHOP	0	625	734 (707-761) ⁶	24 (22-25) ⁶	-	-	13.1
	Germany				694 (670-718) ⁶	20 (19-22) ⁶	-	-	11.5

Age	Country	Study	Breastfeeding ¹ (%)	N	Mean energy (SD) ² (kcal/d) ³	Mean protein (SD) ² (g/d)	P5 th of protein intake ³	P2.5 th of protein intake ⁴	Mean E% as protein ²
	Italy				817 (796-838) ⁵	28 (27-28) ⁶		-	13.7
	Poland				812 (783-840) ⁵	25 (24-26) ⁶	-	-	12.3
	Spain				854 (831-878) ⁵	30 (29-32) ⁶	-	-	14.1
	Finland	STRIP	0	215	842 (148)	25.0 (6)	15.1	13.2	11.9
	UK	ALSPAC	0	618 m	840 (173)	29.0 (9.0)	14.1	11.4	13.8
				513 f	784 (156)	27.0 (8.0)	13.8	11.3	13.8
9 mo	Belgium	CHOP	0	617	770 (739-801) ⁵	24 (22-25) ⁶	-	-	13.5
	Germany				708 (684-731) ⁵	20 (19-22) ⁶	-	-	11.9
	Italy				850 (826-874) ⁵	28 (27-28) ⁶	-	-	14.1
	Poland				859 (850-892) ⁵	25 (24-26) ⁶	-	-	12.6
	Spain				872 (846-899) ⁵	30 (29-32) ⁶	-	-	14.7
	Iceland	1995–1996	37	80	760 (678-859) ⁵	28.0 (21.7-35.3) ⁵	-	-	14.4 ⁷
		2005	41	154	754 (629-859) ⁵	22.7 (17.8-27.6) ⁵	-	-	11.9 ⁷
	Netherlands	TNO	-	333	970 (175)	28.8 (6.2)	18.6	16.6	11.9
	Germany	DONALD	17	332	759 (122)	22.4 (5.3)	13.6	12.0	11.8
10-12 mo	France	National	0	63	826 (160)	30.0 (11)	11.8	-	14.4

350 ¹ Either exclusive or partial breastfeeding; ² Unless otherwise noted; ³ Calculated from the mean and the SD assuming a normal distribution of intake; ⁴ Calculated only for sample sizes n≥180; ⁵
351 Median (interquartile range); ⁶ Mean (95% confidence interval); ⁷ Median; - = not reported or not available; m= males; f = females.

352 **Table 5:** Methods for dietary assessment, estimation of breast milk intake and food composition databases used in published studies

Country	Study	Age (months)	Publications	Dietary assessment	Breast milk intake
France	National	6, 10-12	Fantino & Gourmet, 2008	3-d weighed-DR	
Germany	DONALD	3, 6	Hilbig, 2005	3-d weighed-DR	Measured by 'test-weighing' the infant before and after each breast milk meal
UK	ALSPAC	4	Noble and Emmett, 2006	24-h recall	Duration of each breastfeeding was used to estimate the volume of milk consumed; a feed lasting ≥ 10 min was assumed to be 125 mL, or a proportion of this if the feed was of shorter duration (i.e. 12.5 mL for 1 min)
UK	Southampton	6	Marriott et al., 2008	4-d weighed-DR	Estimated using an algorithm based on length of suckling derived from published intake data
5 EU countries	CHOP	6-12	Damianidi et al., 2006	3-d weighed-DR	-
Finland	STRIP	8	Lagström et al., 1997	3-d DR	-
UK	ALSPAC	8	Noble et al., 2001	3-d DR	-
Iceland	1995–1996 2005	9	Thorisdottir et al., 2013	2-d or 3-d weighed DR	Measured by 'test-weighing' the infant before and after each breast milk meal
Netherlands	TNO	9	de Boer et al., 2006	2-d DR	NR
Germany	DONALD	9	Schwartz et al., 2010	3-d weighed-DR	Measured by 'test-weighing' the infant before and after each breast milk meal

353 DR = Dietary records; NR = not reported.

354 **Table 6:** Mean energy and protein intake in infants living in Europe from the EFSA Comprehensive European Food Consumption Database

Age	Breastfeeding ¹	Country	Survey	Dietary assessment	N	Mean energy intake (kcal/d)	Mean protein intake (g/d)	P5 th of protein intake (g/d)	P2.5 th of protein intake (g/d)	Mean E% as protein
4- <6 mo	Yes	Bulgaria	NUTRICHILD	24-h recall, 3 d ^{2,5}	64	679	11.2	6.9	-	7
		Denmark	IAT 2006_07	7-d DR ^{3,6}	26	746	15.9	-	-	9
		UK	DNSIYC_2011	4-d DR ^{4,7}	27	686	15.0	-	-	9
	No	Bulgaria	NUTRICHILD	24-h recall, 3 d	88	639	16.3	9.7	-	10
		Denmark	IAT 2006_07	7-d DR	12	745	18.2	-	-	10
		UK	DNSIYC_2011	4-d DR	49	639	15.9	-	-	10
≥6-12mo	Yes	Bulgaria	NUTRICHILD	24-h recall, 3 d ^{2,5}	89	905	22.2	8.3	-	10
		Denmark	IAT 2006_07	7-d DR ^{3,6}	315	832	23.0	10.1	8.9	11
		UK	DNSIYC_2011	4-d DR ^{4,7}	264	804	24.5	13.0	12.0	12
	No	Bulgaria	NUTRICHILD	24-h recall, 3 d	343	859	27.4	14.1	11.9	13
		Denmark	IAT 2006_07	7-d DR	473	933	30.0	16.2	14.7	13
		UK	DNSIYC_2011	4-d DR	1029	790	25.2	13.0	11.7	13

355 - = P5th and P2.5th of protein intake are only provided for study groups with a sample size ≥ 60 and 180 individuals, respectively.

356 DR = dietary records; ¹ Either exclusive or partial breastfeeding; ² Method to estimate volume of breast milk intake per feeding occasion not reported, assumed to be 130
 357 mL per feeding occasion; ³ Volume of breast milk per feeding occasion calculated as follows (Dewey et al., 1984): 130 mL per breastfeeding if the infant was breastfed 6
 358 times or more per day; 89 mL per breastfeeding if the infant was breastfed 3-5 times per day; 53 mL per breastfeeding if the infant was breastfed 1-2 times per day; ⁴
 359 Volume of breast milk per feeding occasion calculated based on the time for each feed, at 13.5g/min with a maximum of 135g per feed for infants aged 4 to 7 months and
 360 at 10g/min with a maximum of 100g per feed for infants aged 8 to 12 months; ⁵ Breast milk assumed to contain 70.0 kcal/100mL; 1.0 g of protein/100mL and 1.4 g of
 361 protein/100 kcal; ⁶ Breast milk assumed to contain 71.0 kcal/100mL; 1.3 g of protein/100mL and 1.8 g of protein/100 kcal; ⁷ Breast milk assumed to contain 67.0
 362 kcal/100mL; 1.3 g of protein/100mL and 1.9 g of protein/100 kcal

363

364
365

Table 7: Mean energy and protein intake from formula and complementary food in infants from surveys in the EFSA Comprehensive European Food Consumption Database

Age	Breastfeeding ¹	Country	N	Food group	Mean energy intake (kcal/d)	Mean protein intake (g/d)	Mean protein intake (g/100kcal) ²
4-<6 mo	Yes	Bulgaria	64	CF	210	4.1	-
		Denmark	26	CF	255	6.0	-
		UK	27	CF	211	3.7	-
	No	Bulgaria	88	Formula ³	298	6.5	2.18
				CF	341	9.8	-
		Denmark	12	Formula	495	12.0	2.42
				CF	250	6.2	-
		UK	49	Formula	493	10.9	2.21
CF	146	6.0	-				
≥6-12mo	Yes	Bulgaria	89	CF	554	17.3	-
		Denmark	315	CF	540	17.0	-
		UK	264	CF	405	16.6	-
	No	Bulgaria	343	Formula	91	2.2	2.42
				CF	749	24.8	-
		Denmark	473	Formula	225	6.1	2.71
				CF	708	23.9	-
		UK	1029	Formula	349	7.4	2.12
CF	441	17.6	-				

366 CF = Complementary food; ¹ Either exclusive or partial breastfeeding; ² Calculated from mean energy and protein intake from formula; ³ Any (infant and follow-on) formula

367

368 **Table 8:** Mean energy and protein intake by food group in non-breastfed infants aged 4 to 6 months

Food group	Bulgaria (n=88)		Denmark (n=12)		UK (n=49)	
	Mean energy (kcal/d)	Mean protein (g/d)	Mean energy (kcal/d)	Mean protein (g/d)	Mean energy (kcal/d)	Mean protein (g/d)
Animal and vegetable fats and oils	8.6	0.0	27.4	0.0	1.0	0.0
Composite food (including frozen products)	0.2	0.0	0.0	0.0	0.4	0.0
Eggs and egg products	3.1	0.2	0.0	0.0	0.4	0.0
Fish and other seafood (including amphibians, reptiles, snails and insects)	0.0	0.0	0.4	0.1	0.7	0.1
Food for infants and small children	129.5	2.4	105.5	3.0	95.1	3.0
Fruit and fruit products	7.9	0.1	14.7	0.1	11.5	0.1
Fruit and vegetable juices	15.6	0.1	0.0	0.0	0.7	0.0
Grains and grain-based products	26.6	0.6	47.5	1.1	4.7	0.1
Herbs, spices and condiments	0.0	0.0	0.0	0.0	0.3	0.0
Infant formula and follow-up formula	298.0	6.5	495.2	12.0	493.4	10.9
Legumes, nuts and oilseeds	0.4	0.0	3.7	0.3	1.9	0.2
Meat and meat products (including edible offal)	7.3	0.6	1.0	0.1	2.3	0.4
Milk and dairy products	98.1	5.5	11.1	0.7	13.5	0.8
Snacks, desserts, and other foods	2.1	0.1	0.0	0.0	3.9	0.1
Starchy roots and tubers	9.0	0.2	27.8	0.6	5.0	0.1
Sugar and confectionary	31.6	0.0	4.7	0.0	0.7	0.0
Vegetables and vegetable products (including fungi)	1.2	0.0	5.9	0.3	3.7	0.2

369

370 **Table 9:** Mean energy and protein intake by food group in non-breastfed infants aged 6 to 12 months

Food group	Bulgaria (n=343)		Denmark (n=473)		UK (n=1029)	
	Mean energy (kcal/d)	Mean protein (g/d)	Mean energy (kcal/d)	Mean protein (g/d)	Mean energy (kcal/d)	Mean protein (g/d)
Animal and vegetable fats and oils	70.4	0.0	87.0	0.0	15.8	0.0
Composite food (including frozen products)	1.3	0.1	0.0	0.0	19.2	0.9
Eggs and egg products	13.0	0.9	2.5	0.2	2.9	0.2
Fish and other seafood (including amphibians, reptiles, snails and insects)	1.3	0.2	7.9	1.0	7.2	0.8
Food for infants and small children	99.7	2.4	68.9	1.9	121.4	4.0
Fruit and fruit products	35.6	0.3	62.4	0.7	33.2	0.4
Fruit and vegetable juices	24.8	0.2	4.7	0.0	1.7	0.0
Grains and grain-based products	183.1	4.3	186.3	5.3	78.7	2.4
Herbs, spices and condiments	0.9	0.1	2.0	0.0	3.9	0.1
Infant formula and follow-up formula	90.8	2.2	225.1	6.1	349.0	7.4
Legumes, nuts and oilseeds	6.5	0.4	5.9	0.4	5.5	0.4
Meat and meat products (including edible offal)	48.3	5.2	49.4	3.7	21.9	2.9
Milk and dairy products	166.5	9.2	154.5	8.7	79.3	4.4
Snacks, desserts, and other foods	7.7	0.1	12.0	0.4	12.1	0.2
Starchy roots and tubers	24.0	0.6	34.8	0.8	20.9	0.4
Sugar and confectionary	50.6	0.0	17.1	0.0	5.9	0.1
Vegetables and vegetable products (including fungi)	14.3	0.6	12.5	0.6	10.4	0.5

371

372 Mean protein intakes from all sources were beyond the PRI for protein in all surveys from the EFSA
373 Comprehensive Database for both breastfed and formula-fed infants aged 6 to 12 months (**Table 6**).
374 Mean protein intake from all sources were also reported in published studies which accurately
375 estimated breast milk intake (by weighing the infant before and after each breast milk meal) and/or
376 which used more accurate methods for dietary assessment (3 or 4-day weighted dietary records). At 4
377 months of age, the lowest mean protein intake (12 g per day) were reported for infants exclusively
378 breastfed in the ALSPAC cohort. Mean protein intakes were slightly higher (13 g per day) in breastfed
379 infants who had already received some solid food. The lowest mean protein intake (16.3 g per day)
380 for infants aged 6 months or older was reported in a German cohort (DONALD study), in which the
381 proportion of breastfed infants was the highest (48%) among all the studies available (**Table 4**).
382 Mean protein intakes from all sources were beyond the PRI for protein for infants aged 6 to 12
383 months in all the studies.

384 In the dietary surveys for which data on (exclusively or partially) breastfed infants and formula-fed
385 infants was available separately (**Table 6**), mean protein intakes were systematically higher in
386 formula-fed infants than in breastfed infants, as previously reported by others (Heinig et al., 1993).
387 Breast milk was assumed to contain from 1.4 to 1.9 g of protein/100 kcal, depending on the survey.
388 The lower mean protein intake reported for Bulgarian infants could be explained in part by the
389 assumed lower protein content in breast milk (1.4 g/100 kcal). The protein content of formula ranged
390 from 2.1 to 2.7 g of protein/100 kcal, depending on the survey and age category (**Table 7**). This is
391 higher than the minimum protein content allowed by EU legislation (Directive 2006/141/EC and
392 Commission delegated Regulation (EU) 2016/127) for (infant and follow-on) formula manufactured
393 from intact cows' or goats' milk proteins (1.8 g/100 kcal).

394 In breastfed infants, mean protein intake from CF ranged between 3.7 and 6.0 g/day in infants aged 4
395 to <6 months, and were already well beyond the PRI for protein in infants aged 6-12 months (about
396 17 g/day). In formula-fed infants aged 4 to 6 months, mean protein intake from formula ranged from
397 6.5 to 12.0 g per day, whilst mean protein intakes from CF were about 6 g/day. In Bulgaria, where
398 mean protein intake from formula was the lowest (6.5 g/day), mean protein intake from CF was much
399 higher (9.8 g/day), mostly coming from cow's milk and dairy products other than formula (**Table 8**).
400 This is due to a replacement of IF with cow's milk (rather than with FOF) at the time of the
401 introduction of CF. In formula-fed infants aged 6 to 12 months, mean protein intakes from CF were at
402 or beyond the PRI in all countries. The contribution of formula to total protein intake varied widely,
403 being lower in countries (Bulgaria and Denmark) with the highest protein intake from cow's milk and
404 dairy products and from meat and meat products (**Table 9**).

405 Whenever the data available allowed doing so, the 5th and the 2.5th percentiles (P5th and P2.5th,
406 respectively) of protein intake were calculated by assuming a normal distribution of protein intake
407 data (**Table 4**) or extracted from individual data (**Table 6**). Otherwise, IQRs or 95% CI were
408 considered (**Table 4**). The Panel notes that the P5th and P2.5th of total protein intake in non-breastfed
409 infants aged 6 to 12 months was around or above the PRI for protein for that age group in all the
410 studies (**Table 4**) and surveys (**Table 6**) available.

411 **3.4. Contribution that a FOF with a protein content of at least 1.6 g/100** 412 **kcal could make towards protein requirements in the target** 413 **population**

414 Consumption of a FOF with a protein content of about 1.6 g/100 kcal would provide about 9 g of
415 protein per day in the first months of complementary feeding (assuming an intake of about 500 mL
416 per day) and about 4.5 g of protein per day (assuming an intake of about 250 mL per day) by the end
417 of the first year of life. This is about 1 g and 0.5 g of protein less than the estimated intake from a
418 formula containing a minimum of 1.8 g/100 kcal, as currently authorised. The Panel notes, however,
419 that the protein content of (infant and follow-on) formula in the European surveys available (from 2.1
420 to 2.7 g of protein/100 kcal, **Table 7**) was higher than the minimum authorised.

421 Using individual data from the three surveys which were available in the EFSA Comprehensive Food
422 Consumption Database, total protein intake in non-breastfed infants aged 6 to 12 months was
423 recalculated by assuming that: a) all FOF consumed by the infants contained 1.6 g of protein/100
424 kcal; b) the energy content of the individual FOFs did not change; c) protein intake from other sources

425 (IF, CF) did not change. The mean, P5th and P2.5th of total protein intake under these conditions are
 426 shown in **Table 10**.

427 **Table 10:** Protein intake in European non-breastfed infants aged 6 to 12 months, assuming a
 428 protein content of FOF of 1.6 g/100kcal

Country	N	Mean energy intake (kcal/d)	Mean protein intake (g/d)	P5 th of protein intake (g/d)	P2.5 th of protein intake (g/d)
Bulgaria	343	859	27.2	13.7	11.9
Denmark	473	933	29.4	15.8	13.8
UK	1029	790	24.4	12.6	11.2

429

430 As expected, total protein intakes resulting from the consumption of FOF with a protein content of 1.6
 431 g/100kcal would be lower than those reported in the original surveys (**Table 6**). The Panel notes,
 432 however, that the P5th and P2.5th of total protein intake would remain at about or above the PRI for
 433 protein for infants aged 6 to 12 months.

434 3.5. Application submitted by the food business operator

435 The applicant provided two human intervention studies aiming to investigate whether protein content
 436 in formula to be fed from three to 12 months of age that is closer to the protein content of breast milk
 437 during that feeding period would lead to growth rates more in line to those of breast-fed infants, as
 438 compared to infants fed a "standard" formula.

439 3.5.1. Composition of the formulae used in the two human intervention studies

440 The formulae investigated in the US and Chile studies contain a minimum of 1.61 g protein per 100
 441 kcal, based on a calculation of total nitrogen x 6.25. The protein source is based on intact proteins
 442 derived from skimmed milk and a proprietary preparation of demineralised whey. The demineralised
 443 whey is obtained from modified caseino-glyco-macro-peptide (CGMP)-reduced sweet whey produced
 444 using a patented process (Patent No PCT/EP1998/003176). The whey preparation used in the formula
 445 has a CGMP content which is reduced by at least 85%. The whey protein-to-casein ratio of the final
 446 product is 60:40. The applicant indicated that the protein sources have been used in other FOF
 447 currently marketed by the applicant, and that the use of CGMP-reduced sweet whey has allowed for a
 448 lower protein content of the FOF, while still meeting the requirements of Directive 2006/141/EC with
 449 respect to the amino acid pattern. The energy content and the amount of carbohydrates, fat, vitamins
 450 and minerals also comply with the compositional requirements laid down in Directive 2006/141/EC.

451 The macronutrient composition of the intervention and control formulae used in the US and the Chile
 452 studies are outlined in **Table 11**.

453 **Table 11:** Macronutrient composition of study formulae in comparison to the compositional
 454 requirements for FOF manufactured from cow's or goat's-milk proteins as laid down in
 455 Directive 2006/141/EC

	Unit	Directive 2006/141/EC	US study		Chile study	
			Intervention	Control	Intervention	Control
Energy	kcal/100 ml	60-70	67.2	64.6	62.8	65.6
Protein	g/100 kcal	1.8-3.5	1.61	2.15	1.65	2.70
Fat	g/100 kcal	4.0-6.0	5.46	5.21	5.30	5.03
Carbohydrates	g/100 kcal	9.0-14.0	11.10	11.13	11.41	10.98
Cyst(e)ine	mg/100 kcal	38	28	38	28	46
Histidine	mg/100 kcal	40	40	49	39	64
Isoleucine	mg/100 kcal	90	95	125	100	166
Leucine	mg/100 kcal	166	166	222	180	298
Lysine	mg/100 kcal	113	132	185	142	234
Methionine	mg/100 kcal	23	33	46	42	69
Phenylalanine	mg/100 kcal	83	103	88	108	179

Threonine	mg/100 kcal	77	94	141	84	137
Tryptophan	mg/100 kcal	32	31	31	34	57
Tyrosine	mg/100 kcal	76	52	68	69	113
Valine	mg/100 kcal	88	94	137	102	168
Methionine + Cyst(e)ine	mg/100 kcal	61 ^(a)	61	84	70	115
ratio			1.2	1.2	1.5	1.5
Tyrosine + Phenylalanine	mg/100 kcal	159 ^(b)	155	156	177	292
ratio			0.5	0.8	0.6	0.6

456 (a): The concentrations of cyst(e)ine and methionine may be added together if the methionine:cyst(e)ine-ratio is not >3

457 (b): The concentrations of tyrosine and phenylalanine may be added together if the tyrosine:phenylalanine-ratio is not >2

458

459 The applicant states that the tyrosine and phenylalanine content (calculated as sum) and tryptophan
 460 content in the intervention and control formulae of the US study were slightly lower than required by
 461 Directive 2006/141/EC, and that the histidine content in the intervention formula of the Chile study
 462 was slightly lower than required by Directive 2006/141/EC, but that the final marketed product will
 463 comply with the specifications laid down in the Directive.

464 In the Chile study the intervention formula also contained 2×10^7 colony forming units (CFU)
 465 *Bifidobacterium lactis* (CNCM I-3446) and 2×10^7 CFU *Lactobacillus rhamnosus* (CGMCC 1.3724) per
 466 gram of powder formula, while the control formula did not contain these bacteria.

467 The intervention formulae in both studies had a whey protein-to-casein ratio of 60:40.

468 3.5.2. Human intervention studies

469 The two randomised, double-blind, controlled intervention studies were conducted in Chile (Inostroza,
 470 et al., 2014) and in the US (Ziegler et al., 2015). These studies assessed growth rates in healthy term
 471 infants who consumed (low-protein, intervention) formulae with protein contents of 1.61 g/100 kcal (n
 472 = 97) (US study) and 1.65 g/100 kcal (n = 89) (Chile study) from three months of age onwards,
 473 against those of infants who consumed (control) formulae with protein contents of 2.15 g/100 kcal (n
 474 = 97) and 2.70 g/100 kcal (n = 87), respectively, and against those of a breastfed reference group
 475 (n= 76 and n = 112, respectively). In the Chile study, only infants from overweight and obese
 476 mothers were recruited.

477 In both studies, the primary outcome was weight gain between 3 and 6 months of age. Secondary
 478 outcomes included, amongst others, weight gain at time points beyond 6 months of age, weight
 479 changes, changes in length and head circumference, and changes in serum albumin and blood urea
 480 nitrogen (BUN). Adverse events were registered.

481 Statistical analyses were conducted in completers and per protocol (PP) in both studies. In the US
 482 study, 10 infants in the intervention group, 10 infants in the control group and 7 infants in the
 483 breastfed reference group discontinued the study. The numbers in the Chile study were 23, 11 and
 484 11, respectively. Reasons for withdrawal were provided.

485 Despite the original protocols foreseeing exclusive formula or breastfeeding up to the age of 6 months
 486 and the introduction of CF thereafter (control formula was allowed from 6 to 12 months to the
 487 breastfed reference groups, if desired), small amounts of CF were provided to some infants from
 488 around 4 months of age onwards. In the US study, a total of 9 infants consumed >4 teaspoons of CF
 489 per day before the age of 6 months and were excluded from the PP analysis. In the Chile study, CF in
 490 amounts > 4 teaspoons per day were introduced before 6 months of age in 66 infants (28 in the
 491 intervention, 24 in the control and 14 in the breastfed reference group), who were not excluded from
 492 the statistical analysis.

493 The information provided in these studies did not allow the calculation of energy and protein intake
 494 from CF at any time point, and thus of the total energy and protein intake at time points in which
 495 infants consumed CF (4-12 months in the Chile study, 6-12 months in the US study).

496 Mean daily energy and protein intake from formula at different time points in the US and Chile studies
 497 are given in Table 12.

498
499**Table 12:** Mean daily energy and protein intake from formula at different time-points in the US and Chile studies

US Study				
	n	Mean (SD) volume intake (mL/day)	Mean (SD) energy intake (kcal/day)	Mean (SD) protein intake (g/day)
4 months				
Low-protein formula	83	905 (216)	581 (139)	14.6 (3.5)
Control formula	85	894 (180)	601 (121)	19.2 (3.9)
6 months				
Low-protein formula	83	917 (232)	589 (149)	14.8 (3.7)
Control formula	84	902 (184)	606 (124)	19.4 (4.0)
8 months				
Low-protein formula	80	850 (208)	546 (133)	13.7 (3.3)
Control formula	82	857 (179)	606 (124)	18.4 (3.8)
12 months				
Low-protein formula	76	719 (239)	462 (153)	11.6 (3.8)
Control formula	78	725 (241)	487 (162)	15.6 (5.2)
Chile study				
	n	Mean (SD) volume intake (mL/day)	Mean (SD) energy intake (kcal/day)	Mean (SD) protein intake (g/day)
4 months				
Low-protein formula	75	820 (268)	515 (168)	13.5 (4.4)
Control formula	80	868 (228)	569 (150)	23.4 (6.2)
6 months				
Low-protein formula	62	980 (248)	615 (156)	16.2 (4.1)
Control formula	74	957 (172)	628 (113)	25.8 (4.6)
9 months				
Low-protein formula	55	896 (256)	563 (161)	14.8 (4.2)
Control formula	64	869 (242)	570 (159)	23.5 (6.5)
12 months				
Low-protein formula	47	854 (324)	536 (203)	14.1 (5.3)
Control formula	63	747 (217)	490 (142)	20.2 (5.9)

500

501 The mean volume of formula consumed by infants did not differ significantly between the low protein
 502 and the control groups at any time point in any of the studies. In this context, mean energy intake
 503 from formula was comparable between the two formula groups whilst mean protein intake was
 504 systematically higher in the control formula vs. the low-protein formula groups in both studies, as per
 505 study design. The Panel notes that the mean volume intake of formula in the low-protein formula and
 506 control formula groups in both studies at 9 and 12 months of age was high.

507 The results of anthropometric measurements in the low-protein formula, control formula and
 508 breastfed reference groups in both studies are given in Appendices A-D. In both studies, weight gain

509 (in g/day) was somewhat lower in the infants consuming the low protein formula than in infants
510 consuming the control formula for the time period 3 to 6 months and 6 to 12 months of age, but this
511 difference only reached statistical significance for weight gain between 3 and 6 months in the Chile
512 study. Other anthropometric measures (i.e. weight, length and head circumference at different time
513 points in absolute values and as change from baseline) generally followed this pattern in both studies.

514 In the US study, both formula groups showed statistically significantly higher weight gain and higher
515 weight and length in absolute values at different time points as compared to the breastfed reference
516 group. In the Chile study, the low-protein formula group and the breastfed reference group did not
517 differ significantly in weight gain, weight and length. The concentrations of serum albumin and BUN
518 remained within the normal range in all groups during the intervention in both studies. Reported
519 adverse events were similar in the intervention and control groups.

520 The Panel notes that, in both studies, no differences in growth patterns were observed between
521 infants in the control *vs.* the low-protein formulae, including the time period of 3 to 6 months of age
522 when the formula was fed almost exclusively. In the US study, growth was higher in the low-protein
523 groups as compared to the breastfed reference group, whereas in the Chile study, infants in the low-
524 protein formula group had a similar growth pattern to breastfed infants. The control formula used in
525 these studies contained 0.35 g/100 kcal (US study) and 0.90 g/100 kcal (Chile study) more protein
526 than the current minimum requirement for protein content of a FOF (1.8 g/100 kcal). In both studies,
527 at all-time points, the difference in mean protein intake from formula between the control formula
528 group and the low-protein formula group was 4 g/day or greater.

529 The Panel also notes that the studies submitted were not specifically designed to meet the regulatory
530 definitions for either IF or FOF laid down in Regulation (EU) No 609/2013¹⁰, and that the information
531 provided in relation to the type and amount of CF was not sufficient to calculate total energy and
532 protein intake, nor the relative contribution of formulae and CF to total energy and protein intake.
533 Therefore, the Panel considers that these studies do not provide, on their own, sufficient information
534 to conclude on the safety and suitability of a FOF with a protein content of 1.6 g protein/100kcal.

535 **3.6. Comparison between the human intervention studies provided and** 536 **European dietary surveys with respect to mean energy and protein** 537 **intake from formula and complementary food in the target** 538 **population**

539 In the European surveys which allowed calculation of mean energy and protein intake from both
540 (infant and follow-on) formula and CF in formula-fed infants, the protein content of formula was
541 between 2.1 and 2.7 g/100kcal (Table 7). The lower end is close to the protein content of the control
542 formula used in the US study (2.15 g/100kcal) and the upper end is close to the protein content of the
543 control formula used in the Chile study (2.70g/kcal). Mean energy and protein intakes from formula
544 were, however, lower in infants aged 4-<6 months in the European surveys than in infants at 4
545 months of age in the two formula groups (low protein and control) in both intervention studies (US
546 and Chile studies). In infants aged 6-12 months, mean energy and protein intakes from formula in the
547 European surveys were about half (or lower) than in the US and Chile studies. This suggests that the
548 contribution of formula (*vs.* CF) to total protein intake in the target population (infants at the time of
549 the introduction of complementary feeding and up to 12 months of age) may be lower in Europe than
550 in the intervention studies provided. Therefore, the impact on total protein intake of lowering the
551 protein content of a follow-on formula to about 1.6 g/100 kcal would also be lower in Europe.
552 However, direct comparisons regarding total energy and protein intake and energy and protein intake
553 from CF between the European surveys and the intervention studies provided cannot be made.

¹⁰ Regulation (EU) No 609/2013 of the European Parliament and of the Council of 12 June 2013 on food intended for infants and young children, food for special medical purposes, and total diet replacement for weight control and repealing Council Directive 92/52/EEC, Commission Directives 96/8/EC, 1999/21/EC, 2006/125/EC and 2006/141/EC, Directive 2009/39/EC of the European Parliament and of the Council and Commission Regulations (EC) No 41/2009 and (EC) No 953/2009, OJ L 181, 29.6.2013, p. 35–56

554 **4. Conclusions**

555 **4.1. On the safety and suitability for use by infants of FOF with a protein** 556 **content of at least 1.6 g/100 kcal from intact cow's milk protein** 557 **otherwise complying with the requirements of relevant EU** 558 **legislation**

559 The Panel considers that the two intervention studies provided by the applicant do not provide, on
560 their own, sufficient information to conclude on the safety and suitability of FOF with a protein content
561 of 1.6 g protein/100kcal.

562 The Panel notes, however, that:

- 563 a) the protein content of human milk tends to decrease with feeding time to about 1.1-1.4
564 g/100kcal by the end of the fourth month of lactation, remaining fairly stable thereafter,
- 565 b) that P5th and P2.5th of protein intake from all sources (breast milk, formula and CF) in
566 European infants between 6 and 12 months of age are at or above the PRI for protein for
567 that age group.
- 568 c) that P5th and P2.5th of protein intake from all sources (formula and CF) in European infants
569 between 6 and 12 months of age who are not breastfed would remain at or above the PRI
570 for protein for that age group by assuming a protein content of 1.6 g/100kcal in all FOF.
- 571 d) the two human intervention studies provided by the applicant did not show an adverse
572 impact on growth resulting from the use of a formula containing about 1.6 g of
573 protein/100kcal as compared to control formulae containing 2.15 or 2.70 g of protein/100
574 kcal or the breastfed reference group.

575 Therefore, the Panel concludes that the use of FOF with a protein content of at least 1.6 g/100
576 kcal from intact cow's milk protein otherwise complying with the requirements of relevant EU
577 legislation is safe and suitable for infants living in Europe with access to complementary foods of
578 a sufficient quality. This conclusion does not apply to IF.

579 **4.2. On the safety and suitability for use by infants of FOF with a protein** 580 **content of at least 1.6 g/100 kcal from goat's milk intact protein,** 581 **soy protein isolates or protein hydrolysates otherwise complying** 582 **with the requirements of relevant EU legislation**

583 On the basis of:

- 584 a) a previous evaluation by the Panel on the safety and suitability of goat's milk protein as a
585 source of protein in IF and FOF (EFSA NDA Panel, 2012b), and
- 586 b) the Panel's conclusions regarding the safety and suitability of FOF with a protein content of at
587 least 1.6 g/100 kcal from intact cow's milk protein otherwise complying with the requirements
588 of relevant EU legislation (section 4.1),

589 the Panel concludes that the use of FOF with a protein content of at least 1.6 g/100 kcal from intact
590 goat's milk protein otherwise complying with the requirements of relevant EU legislation is safe and
591 suitable for infants living in Europe with access to complementary foods of a sufficient quality. This
592 conclusion does not apply to IF.

593 The Panel considers, however, that the safety and suitability of each FOF (and IF) manufactured from
594 protein hydrolysates have to be established by clinical evaluation in the target population (EFSA NDA
595 Panel, 2014). The Panel also considers that, given the higher minimum protein requirements
596 established for FOF (and IF) manufactured from soy protein isolates (i.e. 2.25 g/100 kcal) and the
597 lack of data available on the use of FOF from soy protein isolates in the target population, additional
598 studies are required to establish the safety and suitability of FOF manufactured from soy protein
599 isolates with a protein content of at least 1.6 g/100 kcal. Therefore, the Panel concludes that the
600 safety and suitability of FOF with a protein content of at least 1.6 g/100 kcal manufactured from either

601 protein hydrolysates or soy protein isolates cannot be established with the available data. The same
602 conclusion applies to IF.

DRAFT

603 **Documentation provided to EFSA**

604 Application for the placing on the market of a follow-on formula with a new minimum protein content
605 of 1.61 gram of protein per 100 kcal. April 2016. Nestlé Nutrition.

606 **References**

- 607 AAP (American Academy of Pediatrics), 1988. Clinical testing of infant formulas with respect to
608 nutritional suitability for term infants. Report prepared under FDA contract 223-86-2117, available
609 online: [http://www.fda.gov/Food/GuidanceRegulation/GuidanceDocumentsRegulatoryInformation/](http://www.fda.gov/Food/GuidanceRegulation/GuidanceDocumentsRegulatoryInformation/InfantFormula/ucm170649.htm)
610 [InfantFormula/ucm170649.htm](http://www.fda.gov/Food/GuidanceRegulation/GuidanceDocumentsRegulatoryInformation/InfantFormula/ucm170649.htm).
- 611 Aggett PJ, Agostini C, Goulet O, Hernell O, Koletzko B, Lafeber HL, Michaelsen KF, Rigo J, Weaver LT
612 and ESPGHAN Committee on Nutrition, 2001. The nutritional and safety assessment of breast milk
613 substitutes and other dietary products for infants: a commentary by the ESPGHAN Committee on
614 Nutrition. *Journal of Pediatric Gastroenterology and Nutrition*, 32, 256-258.
- 615 Allen JC, Keller RP, Archer P, Neville MC, 1991. Studies in human lactation: milk composition and daily
616 secretion rates of macronutrients in the first year of lactation. *American Journal of Clinical Nutrition*
617 54, 69-80.
- 618 Damianidi L, Gruszfeld D, Verduci E, Vecchi F, Xhonneux A, Langhendries JP, Luque V, Theurich MA,
619 Zaragoza-Jordana M, Koletzko B, Grote V, 2016. Protein intake and their nutritional sources during
620 the first 2 years of life: secondary data evaluation from the European Childhood Obesity Project.
621 *European Journal of Clinical Nutrition*, 70, 1291–1297.
- 622 de Boer EJ, Hulshof KFAM, ter Doest D (2006) Voedselconsumptie van jonge peuters [Food
623 consumption of young children]. TNO rapport V6269.
- 624 EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2012a. Scientific Opinion
625 on Dietary Reference Values for protein. *EFSA Journal* 2012;10(2):2557, 66 pp.
626 doi:10.2903/j.efsa.2012.2557
- 627 EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2012b. Scientific Opinion
628 on the suitability of goat milk protein as a source of protein in infant formulae and in follow-on
629 formulae. *EFSA Journal* 2012;10(3):2603, 18 pp. doi:10.2903/j.efsa.2012.2603
- 630 EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2013. Scientific Opinion on
631 nutrient requirements and dietary intake of infants and young children in the European Union.
632 *EFSA Journal* 2013;11(10):3408, 103 pp. doi:10.2903/j.efsa.2013.3408
- 633 EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2014. Scientific Opinion on
634 the essential composition of infant and follow-on formulae. *EFSA Journal* 2014;12(7):3760, 106 pp.
635 doi:10.2903/j.efsa.2014.3760
- 636 Fantino M, Gourmet E., Apports nutritionnels en France en 2005 chez les enfants non allaités âgés
637 de moins de 36 mois (Nutrient intake in 2005 by non-breast fed French children of less than 36
638 months), 2008. *Archives de Pédiatrie* 15,,446–455.
- 639 FAO/WHO/UNU (Food and Agriculture Organization of the United Nations, World Health Organization,
640 United Nations University), 2007. Protein and amino acid requirements in human nutrition. Report
641 of a joint FAO/WHO/UNU expert consultation, WHO Technical Report Series 935.
- 642 Gidrewicz DA and Fenton TR, 2014. A systematic review and meta-analysis of the nutrient content of
643 preterm and term breast milk. *BMC Pediatrics*, 14, 216. doi: 10.1186/1471-2431-14-216
- 644 Hayes NP and Northington R, 2014. Unpublished study report #1. Effect of the level of dietary protein
645 on infant growth and body composition in the first year of life (US study).
- 646 Heinig MJ, Nommsen LA, Peerson JM, Lonnerdal B, Dewey KG, 1993. Energy and protein intakes of
647 breast-fed and formula-fed infants during the first year of life and their association with growth
648 velocity: the DARLING Study. *American Journal of Clinical Nutrition* 58, 152-61.

- 649 Hilbig A and Kersting M, 2006. Effects of age and time on energy and macronutrient intake in German
650 infants and young children: Results of the DONALD study. *Journal of Pediatric Gastroenterology*
651 *and Nutrition* 43, 518-524.
- 652 ICH (International Conference on Harmonisation of Technical Requirements for Registration of
653 Pharmaceuticals for Human Use), 1998. *Statistical Principles for Clinical Trials* E9. 39 pp.
- 654 Inostroza J, Haschke F, Steenhout P, Grathwohl D, Nelson SE and Ziegler EE, 2014. Low-protein
655 formula slows weight gain in infants of overweight mothers. *Journal of Pediatric Gastroenterology*
656 *and Nutrition*, 59, 70-77.
- 657 IoM (Institute of Medicine), 2004. *Infant Formula: Evaluating the Safety of New Ingredients*,
658 Committee on the Evaluation of the Addition of Ingredients New to Infant Formula, Food and
659 Nutrition Board. The National Academies Press, Washington, DC, USA. 10.17226/10935.
- 660 Lagström H, Jokinen E, Seppänen R, 1997. Nutrient intake by young children in a prospective
661 randomized trial of a low-saturated fat, low-cholesterol diet. *Archives of Pediatrics and Adolescent*
662 *Medicine*, 151,181-188.
- 663 Marriott LD, Robinson SM, Poole J, Borland SE, Godfrey KM, Law CM, Inskip HM; Southampton
664 Women's Survey Study Group, 2008. What do babies eat? Evaluation of a food frequency
665 questionnaire to assess the diets of infants aged 6 months. *Public Health Nutrition*, 11, 751-756.
- 666 Michaelsen KF, Skafté L, Badsberg JH, Jørgensen M, 1990. Variation in macronutrients in human bank
667 milk: Influencing factors and implications for human milk banking. *Journal of Pediatric*
668 *Gastroenterology and Nutrition* 11, 229-239.
- 669 Mitoulas LR, Kent JC, Cox DB, Owens RA, Sherriff JL, Hartmann PE, 2002. Variation in fat, lactose and
670 protein in human milk over 24 h and throughout the first year of lactation. *British Journal of*
671 *Nutrition* 88, 29-37.
- 672 Noble S and Emmett P, 2001. Original Communication Food and nutrient intake in a cohort of 8-
673 month-old infants in the south-west of England in 1993. *European Journal of Clinical Nutrition*, 55,
674 698-707.
- 675 Noble S and Emmett P, 2006. Differences in weaning practice, food and nutrient intake between
676 breast- and formula-fed 4-month-old infants in England, 2006. *Journal of Human Nutrition and*
677 *Dietetics*, 19, 303-313.
- 678 Nommsen LA, Lovelady CA, Heinig MJ, Lönnerdal B, Dewey KG, 1991. Determinants of energy,
679 protein, lipid, and lactose concentrations in human milk during the first 12 mo of lactation: the
680 DARLING Study. *American Journal of Clinical Nutrition* 53, 457-65.
- 681 SCF (Scientific Committee on Food), 2003. Report of the Scientific Committee on Food on the revision
682 of essential requirements of infant formulae and follow-on formulae. Available online:
683 http://ec.europa.eu/food/fs/sc/scf/out199_en.pdf
- 684 Schwartz J, Dube K, Alexy U, Kalhoff H, Kersting M, 2010. PUFA and LC-PUFA intake during the first
685 year of life: can dietary practice achieve a guideline diet? *European Journal of Clinical Nutrition*, 64,
686 124-130.
- 687 Thorisdottir B, Gunnarsdottir I, Thorisdottir AV, Palsson GI, Halldorsson TI, Thorsdottir I, 2013.
688 Nutrient intake in infancy and body mass index at six years in two population-based cohorts
689 recruited before and after revision of infant dietary recommendations. *Annals of Nutrition and*
690 *Metabolism*, 63,145-151.
- 691 Yao M, 2014. Unpublished study report #2. Growth in infants of overweight or obese mothers, that
692 receive a low protein formula (Chile study).
- 693 Ziegler EE, Fields DA, Chernausek SD, Steenhout P, Grathwohl D, Jeter JM, Nelson SE and Haschke F,
694 2015. Adequacy of infant formula with protein content of 1.6 g/100 kcal for infants between 3 and
695 12 months. *Journal of Pediatric Gastroenterology and Nutrition*, 61, 596-603.

696 **Abbreviations**

AAP	American Academy of Pediatrics
AR	Average requirement
CF	Complementary food
CGMP	Caseinoglycomacropeptide
DR	Dietary records
E%	Percentage of total energy intake
EC	European Commission
ESPGHAN	European Society for Paediatric Gastroenterology, Hepatology and Nutrition
EFSA	European Food Safety Authority
EU	European Union
FAO	Food and Agriculture Organization of the United Nations
FOF	Follow-on formula
IF	Infant formula
IoM	Institute of Medicine
NDA	Dietetic Products, Nutrition and Allergies
P2.5 th	Percentile 2.5
P5 th	Percentile 5
PRI	Population Reference Intake
SCF	Scientific Committee on Food
US	United States
WHO	World Health Organization

Appendix A – Absolute weight, weight gain and weight change in the “US study”

	n	Intervention mean ± SD	n	Control mean ± SD	n	Breast-fed mean ± SD	Intervention vs control mean difference (95%CI) ^(a)	Intervention vs breast-fed mean difference (95%CI) ^(a)	Control vs breast-fed mean difference (95%CI) ^(a)
Weight gain (g/day)							Weight gain (g/day)		
3-6 months									
Completers	92	20.09 ± 4.62	91	20.65 ± 5.54	109	17.43 ± 4.89	-0.67 (-2.11 to 0.77)	-	-
PP	-	-	-	-	-	-	-0.69 (-2.20 to 0.81)	-	-
6-12 months									
Completers	86	11.54 ± 3.04	87	12.31 ± 3.08	103	10.39 ± 3.13	-0.80 (-1.73 to 0.13)	1.04 (0.12 to 1.95)	1.84 (0.93 to 2.75)
PP	78	11.45 ± 3.00	81	12.22 ± 3.05	97	10.28 ± 3.03	-0.85 (-1.80 to 0.10)	0.96 (0.03 to 1.89)	1.81 (0.90 to 2.73)
Weight (kg)							Weight (g)		
3 months									
Completers	97	5.96 ± 0.70	97	5.91 ± 0.68	112	5.78 ± 0.70	-	-	-
PP	85	5.99 ± 0.68	85	5.89 ± 0.69	105	5.79 ± 0.68	-	-	-
6 months									
Completers	93	7.62 ± 0.84	91	7.62 ± 0.89	109	7.24 ± 0.87	-71.06 (-193.4 to 51.25)	167.86 (47.71 to 288.01) ¹	238.92 (118.28 to 359.57) ²
PP	84	7.68 ± 0.84	85	7.62 ± 0.90	103	7.27 ± 0.89	-71.08 (-199.9 to 57.76)	175.72 (49.63 to 301.81)	246.80 (121.83 to 371.77)
12 months									
Completers	87	9.86 ± 1.12	87	9.97 ± 1.21	104	9.20 ± 1.10	-231.7 (-473.2 to 9.86)	385.61 (148.70 to 622.51) ¹	617.28 (380.58 to 853.98) ²
PP	79	9.89 ± 1.12	81	9.95 ± 1.21	98	9.20 ± 1.12	-239.7 (-484.6 to 5.11)	378.73 (139.63 to 617.83)	618.48 (382.08 to 854.88)
Weight change (kg)							Weight change (kg)		
6-12 months									
Completers	86	2.20 ± 0.58	87	2.35 ± 0.59	103	1.96 ± 0.59	-0.16 (-0.34 to 0.02)	0.21 (0.03 to 0.38) ³	0.37 (0.19 to 0.54) ²
PP	78	2.18 ± 0.57	81	2.33 ± 0.58	97	1.94 ± 0.56	-0.17 (-0.35 to 0.01)	0.19 (0.02 to 0.37)	0.36 (0.19 to 0.53) ²

(a) adjusted for baseline and gender; ¹ p ≤ 0.01; ² p ≤ 0.0001; ³ p < 0.05

697
698
699

Appendix B – Length and head circumference in the “US study”

	n	Intervention mean ± SD	n	Control mean ± SD	n	Breast-fed mean ± SD	Intervention vs control mean difference (95%CI) ^(a)	Intervention vs breast-fed mean difference (95%CI) ^(a)	Control vs breast-fed mean difference (95%CI) ^(a)
Length (cm)							Length (mm)		
3 months									
Completers	97	59.54 ± 2.24	97	59.44 ± 2.06	112	59.62 ± 2.04		-	
PP	85	59.62 ± 2.20	85	59.35 ± 2.03	105	59.61 ± 2.03		-	
6 months									
Completers	94	65.61 ± 2.30	91	65.75 ± 2.34	110	65.11 ± 2.32	-1.43 (-4.51 to 1.65)	6.03 (3.02 to 9.04) ¹	7.46 (4.41 to 10.51) ²
PP	85	65.72 ± 2.35	85	65.63 ± 2.30	104	65.16 ± 2.35	-1.48 (-4.63 to 1.67)	5.79 (2.73 to 8.86) ¹	7.27 (4.19 to 10.35) ²
12 months									
Completers	87	74.52 ± 2.53	87	74.71 ± 2.77	105	73.34 ± 2.69	-2.37 (-7.24 to 2.50)	12.04 (7.28 to 16.80) ²	14.41 (9.63 to 19.19) ²
PP	79	74.58 ± 2.55	81	74.55 ± 2.74	99	73.35 ± 2.74	-2.95 (-8.08 to 2.18)	11.67 (6.68 to 16.65) ²	14.62 (9.64 to 19.61) ²
Head circumference (cm)							Head circumference (mm)		
3 months									
Completers	97	40.27 ± 1.24	96	40.34 ± 1.27	112	40.27 ± 1.23		-	
PP	85	40.26 ± 1.16	84	40.41 ± 1.15	105	40.25 ± 1.26		-	
6 months									
Completers	94	42.99 ± 1.30	91	43.19 ± 1.29	110	42.92 ± 1.26	-0.57 (-2.24 to 1.10)	0.89 (-0.74 to 2.52)	1.46 (-0.19 to 3.11)
PP	85	43.00 ± 1.23	85	43.18 ± 1.25	104	42.88 ± 1.26	-0.09 (-1.67 to 1.49)	1.11 (-0.43 to 2.65)	1.20 (-0.35 to 2.74)
12 months									
Completers	86	46.18 ± 1.39	87	46.47 ± 1.49	104	46.04 ± 1.39	-1.36 (-3.82 to 1.11)	1.76 (-0.64 to 4.17)	3.12 (0.70 to 5.54) ³
PP	79	46.23 ± 1.32	81	46.43 ± 1.44	98	45.98 ± 1.37	-0.43 (-2.73 to 1.86)	2.34 (0.10 to 4.57) ³	2.77 (0.54 to 5.00) ³

^(a) adjusted for baseline and gender; ¹ p < 0.001; ² p < 0.0001; ³ p < 0.05

700

Appendix C – Absolute weight, weight gain and weight change in the “Chile study”

	n	Intervention mean ± SD	n	Control mean ± SD	n	Breast-fed mean ± SD	Intervention vs control mean difference (95%CI) ^(a)	Intervention vs breast-fed mean difference (95%CI) ^(a)	Control vs breast-fed mean difference (95%CI) ^(a)	
Weight gain (g/day)							Weight gain (g/day)			
3-6 months										
Completers	66	18.97 ± 4.19	76	20.74 ± 5.01	65	20.07 ± 5.79	-2.26 (-3.88 to -0.64) ¹	-0.72 (-2.46 to 1.01)	1.54 (-0.13 to 3.21)	
PP	55	19.17 ± 4.16	68	21.02 ± 4.88	57	20.35 ± 5.83	-	-	-	
6-12 months										
Completers	54	10.97 ± 3.05	66	12.13 ± 3.03	61	10.18 ± 3.85	-0.88 (-2.10 to 0.35)	0.77 (-0.50 to 2.05)	1.65 (0.45 to 2.85) ¹	
PP	47	11.05 ± 3.03	60	12.09 ± 3.15	54	9.99 ± 3.94	-0.76 (-2.09 to 0.58)	1.16 (-0.24 to 2.57)	1.92 (0.63 to 3.21) ¹	
Weight (kg)							Weight (g)			
6 months										
Completers	66	8.03 ± 0.67	76	8.17 ± 0.95	65	8.50 ± 1.12	-142.91 (-377.33 to 91.52)	3.75 (-245.49 to 252.99)	146.65 (-93.14 to 386.45)	
PP	55	8.09 ± 0.68	68	8.21 ± 0.97	57	8.52 ± 1.11	-161.08 (-502.24 to 180.08)	71.78 (-286.92 to 430.48)	232.85 (102.29 to 568.00)	
12 months										
Completers	54	10.08 ± 0.86	66	10.36 ± 1.10	61	10.42 ± 1.27	-315.70 (-566.93 to -64.46) ³	132.25 (-129.81 to 394.32)	447.95 (199.13 to 696.77) ²	
PP	47	10.14 ± 0.87	60	10.40 ± 1.13	54	10.43 ± 1.28	-284.55 (-655.77 to 86.68)	273.83 (-112.04 to 659.70)	558.38 (211.97 to 904.79) ¹	
Weight change (kg)							Weight change (kg)			
6-12 months										
Completers	54	2.01 ± 0.55	66	2.20 ± 0.55	61	1.89 ± 0.71	-0.14 (-0.37 to 0.08)	0.11 (-0.12 to 0.35)	0.26 (0.04 to 0.48) ³	
PP	47	2.03 ± 0.54	60	2.20 ± 0.57	54	1.86 ± 0.73	-0.12 (-0.37 to 0.12)	0.19 (-0.07 to 0.44)	0.31 (0.07 to 0.54) ³	

701
702

(a) adjusted for baseline, gender, pre-pregnancy BMI of the mother (as continuous variable), antibiotic use, introduction of complementary food prior to 6 months of age (yes/no) and ethnicity;
¹ p ≤ 0.01; ² p ≤ 0.001; ³ p < 0.05

Appendix D – Length and head circumference in the “Chile study”

	n	Intervention mean ± SD	n	Control mean ± SD	n	Breast-fed mean ± SD	Intervention vs control mean difference (95%CI) ^(a)	Intervention vs breast-fed mean difference (95%CI) ^(a)	Control vs breast-fed mean difference (95%CI) ^(a)
Length (cm)							Length (mm)		
3 months									
Completers	89	59.80 ± 2.01	87	60.34 ± 2.07	76	61.20 ± 1.95	-6.21 (-12.03 to -0.39) ¹	-14.60 (-20.71 to -8.49) ²	-8.39 (-14.47 to -2.30) ³
PP	65	59.84 ± 1.95	73	60.30 ± 2.01	65	61.08 ± 1.93	-6.46 (-12.94 to 0.02)	-14.05 (-20.83 to -7.27) ²	-7.59 (-14.07 to -1.11) ¹
6 months									
Completers	66	66.04 ± 1.78	76	66.54 ± 2.25	65	67.09 ± 2.25	0.08 (-5.36 to 5.53)	1.86 (-3.97 to 7.68)	1.77 (-3.74 to 7.29)
PP	55	66.20 ± 1.82	68	66.56 ± 2.14	57	67.04 ± 2.18	4.12 (-3.75 to 11.99)	8.64 (0.21 to 17.06) ¹	4.51 (-3.26 to 12.28)
12 months									
Completers	54	74.23 ± 1.99	66	74.77 ± 2.34	61	75.37 ± 2.60	-0.75 (-6.58 to 5.08)	1.58 (-4.54 to 7.70)	2.33 (-3.40 to 8.05)
PP	47	74.44 ± 1.97	60	74.81 ± 2.33	54	75.36 ± 2.64	3.19 (-5.33 to 11.71)	8.33 (-0.70 to 17.35)	5.13 (-2.89 to 13.16)
Head circumference (cm)							Head circumference (mm)		
3 months									
Completers	89	40.22 ± 1.06	87	40.50 ± 1.11	76	40.66 ± 1.19	-3.19 (-6.29 to -0.10) ¹	-4.71 (-7.96 to -1.47) ³	-1.52 (-4.75 to 1.72)
PP	65	40.29 ± 0.97	73	40.46 ± 1.07	65	40.74 ± 1.15	-2.52 (-5.92 to 0.89)	-4.97 (-8.53 to -1.40) ³	-2.45 (-5.86 to 0.96)
6 months									
Completers	66	43.10 ± 1.17	76	43.27 ± 1.24	65	43.43 ± 1.36	0.73 (-1.24 to 2.69)	0.79 (-1.28 to 2.86)	0.06 (-1.92 to 2.03)
PP	55	43.22 ± 1.05	68	43.26 ± 1.17	57	43.51 ± 1.31	1.12 (-1.72 to 3.96)	1.76 (-1.25 to 4.77)	0.64 (-2.14 to 3.42)
12 months									
Completers	54	46.02 ± 1.32	66	46.34 ± 1.09	61	46.40 ± 1.36	-0.76 (-2.87 to 1.34)	0.90 (-1.28 to 3.08)	1.67 (-0.38 to 3.71)
PP	47	46.11 ± 1.16	60	46.35 ± 0.97	54	46.50 ± 1.32	-2.19 (-5.32 to 0.93)	0.80 (-2.46 to 4.07)	3.00 (0.11 to 5.88) ¹

703
704

^(a) adjusted for baseline, gender, pre-pregnancy BMI of the mother (as continuous variable), antibiotic use, introduction of complementary food prior to 6 months of age (yes/no) and ethnicity;
¹ p < 0.05; ² p ≤ 0.001; ³ p ≤ 0.01